

***Pseudosperma albobrunneum* sp. nov. from coniferous forests of Pakistan**

SANA JABEEN^{1&2}, ZAINAB², HIRA BASHIR^{2&3}, ABDUL NASIR KHALID²

¹ Department of Botany, Division of Science & Technology,
University of Education, Township, Lahore, Punjab, Pakistan

² Institute of Botany, University of the Punjab,
Quaid-e-Azam Campus-54590, Lahore, Punjab, Pakistan

³ Department of Botany, University of Okara,
Okara 56300, Punjab, Pakistan

* CORRESPONDENCE TO: sanajabeenue@gmail.com; sanajabeen@ue.edu.pk

ABSTRACT—A new species, *Pseudosperma albobrunneum*, is described and illustrated. The specimens are from different areas of Khyber Pakhtunkhwa province, Pakistan. The identification is based on morphological characters, in combination with molecular phylogenetic analysis of sequences of the ITS region of nuclear ribosomal DNA. The species is found distinct morphologically from all closely related taxa, and molecular data confirm its novelty.

KEY WORDS—ectomycorrhizal, Himalayan, moist temperate, taxonomy

Introduction

Inocybaceae Jülich (*Basidiomycota*, *Agaricales*) is a highly diversified family of ectomycorrhizal fungi, comprising seven genera and more than 1000 species (Kobayashi 2009; Matheny & al. 2009, 2012, 2019; Kobayashi & Onishi 2010; Kropf & al. 2010; Bougher & Matheny 2011; Bougher & al. 2012; Kokkonen & Vauras 2012; Fan & Bau 2013, 2014; Braaten & al. 2014; Esteve-Raventós & al. 2015; Jabeen & al. 2016; Farooqi & al. 2017; Naseer & al. 2017; Liu & al. 2018; Ullah & al. 2018; Matheny & Kudzma 2019). The family shows more diversification in temperate regions as comparatively fewer species have been reported from tropical rain forests (Matheny & al. 2003; Matheny &

Bougher 2017). The species have demonstrated ectomycorrhizal associations with as many as 23 families of vascular plants, including low woody shrubs in arctic-alpine habitats making these fungi of high interest (Cripps & al. 2010, Matheny & al. 2019).

Among the 28 species representing different genera in *Inocybaceae* reported from Pakistan (Ahmad & al. 1997; Ilyas & al. 2013; Saba & al. 2015, 2020; Jabeen & al. 2016; Farooqi & al. 2017; Naseer & al. 2017, 2019; Liu & al. 2018; Ullah & al. 2018; Jabeen & Khalid 2020), eight represent *Pseudosperma* (Ahmad & al. 1997; Saba & al. 2015, 2020; Liu & al. 2018; Ullah & al. 2018; Jabeen & Khalid 2020).

The many classification systems proposed for the species in *Inocybaceae* include several clades, sections, subgenera, and genera based on morphological features and molecular phylogenetics (Massee 1904, Heim 1931, Kühner & Romagnesi 1953, Kuyper 1986, Singer 1986, Horak 2005, Matheny & Bougher 2006, Matheny & al. 2009, Larsson & al. 2009, Alvarado & al. 2010, Ryberg & al. 2010, Matheny & Kudzma 2019).

One multigene phylogenetic analysis by Matheny & al. (2019) recognized seven genera within *Inocybaceae*, including two new genera [*Nothocybe* Matheny & K.P.D. Latha, *Pseudosperma* Matheny & Esteve-Rav.], two subgenera elevated to generic rank [*Inosperma* (Kühner) Matheny & Esteve-Rav., *Mallochybe* (Kuyper) Matheny & al.], and three previously described genera [*Auritella* Matheny & Bougher, *Tubariomyces* Esteve-Rav. & Matheny, *Inocybe* (Fr.) Fr. sensu stricto].

Pseudosperma comprises species previously placed in *Inocybe* sect. *Rimosae* (Fr.) Quél. and characterized by cheilocystidia, absence of pleurocystidia, hyaline basidia, adnexed to sinuate lamellae, a fibrillose (rarely squamulose) and rimose pileus surface, a cylindrical stipe with a distinctly pruinose, furfuraceous, or somewhat flocculose apex, stipe context not changing color when bruised, odor often spermatic (reminiscent of green corn or honey) but occasionally nil, and smooth elliptic to (occasionally) indistinctly phaseoliform basidiospores (Matheny & al. 2019). *Pseudosperma* is represented by 70 species in Africa, Asia, Australasia, Europe, North America, and northern South America.

During field surveys for ectomycorrhizal fungi associated with *Pinaceae* in different areas of Khyber Pakhtunkhwa province, Pakistan, several collections were identified through both morphological and nrDNA ITS sequence analyses. They showed characters similar to species now classified under *Pseudosperma*. We could not find a published description that fit these

specimens nor published similar ITS sequences and propose these here as a new species

Material & methods

Samples of basidiomata were collected in four different localities in two administrative divisions of Khyber Pakhtunkhwa province, Pakistan.

Khanian is a small village in the northern Kaghan valley in the Mansehra district, Hazara division, in north-east Khyber Pakhtunkhwa province, immediately south of the main Himalayan range with a typically moist temperate climate and dominated by *Cedrus deodara* (Roxb. ex D. Don) G. Don along with *Abies pindrow* (Royle ex D. Don) Royle and *Pinus wallichiana* A.B. Jacks. (Siddiqui & al. 2013).

The Lower Dir district is located in the Hindu Kush range, Malakand division, in northwestern Khyber Pakhtunkhwa province. The area lies in the valley of the Panjkora river, which originates from the Hindu Kush ranges and joins the Swat River near Chakdara. The climate is dry temperate. *Pinus* spp. and *Quercus oblongata* D. Don [= *Q. incana* Roxb., nom. illeg.] are the dominant tree species (Champion & al. 1965). Among *Pinus*, *P. gerardiana* Wall. ex D. Don was the dominant species. Pure *Q. oblongata* forest was found in the mountain heights (H. Bashir, pers. obs.). Kalam and Mashkun, also situated in the Malakand division, are home to the offshoots of the Hindu Kush range (Hamayun & al. 2003), also dominated by *C. deodara* forests with *Pinus* spp. and *Q. oblongata* (Champion & al. 1965). These areas have a typical dry temperate climate (Stucki & Khan 1999).

Basidiomata were collected and photographed in their natural habitat. Morphological data was recorded from the fresh specimens. Color codes follow Munsell color charts (Munsell 1975). Each collection was dried using a fan heater and preserved in sealed bags. Sections from basidiomata were rehydrated in 5% KOH, stained in Congo red, and observed under a Techno MX4300H compound microscope. Microscopical characters were measured using an ocular micrometer, and drawn with the aid of a camera lucida. The abbreviation (n/m/p) represents 'n' number of basidiospores measured, 'm' number of basidiomata, and 'p' number of collections. Basidiospore dimensions were recorded as (a) b–c (d), where (a) and (d) are the extreme values, and the range b–c contains at least 90% of the calculated values; Q indicates the l/w ratio of the spores, and avQ is the average Q of all measured spores. Measurements of hyphae are given as ranges. The collections examined during this study have been deposited in the herbarium, Department of Botany, University of the Punjab, Quaid-e-Azam Campus, Lahore, Pakistan (LAH).

Genomic DNA extraction and PCR amplification of ITS1, 5.8S, and ITS2 was carried out following procedures by White & al. (1990), Gardes & Bruns (1993), and Bruns (1995). The PCR products were purified and sequenced at Macrogen Inc. (Korea). The newly generated sequences were deposited in GenBank. For phylogenetic study, consensus sequences of ITS regions were generated in the BioEdit software version 7.2.5 (Hall 1999). Sequence homology search was investigated using the BLAST algorithm at <http://www.ncbi.nlm.nih.gov/>. Sequences from the closest

relatives of the Pakistani species were included in the final dataset to reconstruct a phylogeny based on published phylogenies (Larsson & al. 2009, Kropp & al. 2013, Latha & Manimohan 2017, Liu & al. 2018). Sequences from *Auritella* and *Inosperma* were chosen as outgroup (Matheny & al. 2009). GenBank accession numbers are included with each taxon name in the phylogenetic tree. Multiple sequences were aligned using the webPRANK tool at <https://www.ebi.ac.uk/goldman-srv/webprank/>. Maximum likelihood analysis was performed in MEGA version 6 (Tamura & al. 2013) at 1000 bootstrap pseudoreplicates by finding best-fit substitution model with 1000 bootstrap replicates. The phylogeny was inferred by the Maximum Likelihood method based on the General Time Reversible model. A discrete Gamma distribution was used to model evolutionary rate differences among sites (5 categories (+G, parameter = 0.7529)).

Taxonomy

Pseudosperma albobrunneum Jabeen, Zainab, H. Bashir & Khalid, *sp. nov.*

FIGS. 1, 2

MB 840056

Differs from *Pseudosperma dulcamaroides* and *P. sororium* by the smaller basidiomata, white pileus becoming brownish with age, fibrillose stipe, and smooth, elongate to cylindric, sub-reniform basidiospores.

TYPE: Pakistan, Khyber Pakhtunkhwa, Hazara division, Mansehra district, Khanian, 2500 m a.s.l., on soil under *Cedrus deodara*, 5 August 2014, Sana Jabeen SJ102 (Holotype LAH35047; GenBank MG495392).

ETYMOLOGY: The specific epithet refers to the white to brown basidiomata.

PILEUS 1–1.8 cm in diam., conical with a prominent umbo; surface dry, rimose, fibrillose, white (10Y8/4) to brownish (7.5YR6/10); umbo yellowish (5Y8/6) becoming brownish (7.5YR6/10) at maturity; margin incurved when young. LAMELLAE adnexed, moderately close, up to 2 mm deep, golden brown (10YR5/6), with fimbriate edge. STIPE 5.5 × 0.4 cm, central, cylindrical, narrower towards the apex and wider towards the base; surface dry, pruinose towards apex, fibrillose; white (10Y8/4) when young, becoming grayish brown (5R4/1) to brown (7.5YR6/10) upon maturity, context off-white (5Y9/2). ODOR spermatic.

BASIDIOSPORES [100/5/5] (12.7–)13.2–14.4(–16.4) × (5.3–)6.1–6.9 (–8.2) µm, avl × avw = 13.8 × 6.5 µm, Q = (1.90–)1.94–2.20(–2.50), avQ = 2.00, elongate to cylindric, constricted in the center in side view, sub-reniform, smooth, guttulate, brown in 5% KOH. BASIDIA (38.1–)40.3–42.2(–43.7) × (9.2–)9.5–11.8(–12.1) µm, 2–4-spored, clavate, guttulate, hyaline in 5% KOH; clamp connections observed at the base. CHEILOCYSTIDIA (20.2–)25.4–27.4 (–30.3) × 4.4–6.5 µm, avl × avw = 26.4 × 5.5 µm, clavate, connected with basal



FIGURE. 1. *Pseudosperma albobrunneum*. A. LAH35288; B. LAH35047 (holotype). Scale bars = 0.5 cm. Photos by Abdul Nasir Khalid and Sana Jabeen.




FIGURE 2. *Pseudosperma albobrunneum* (holotype, LAH35047). A. Basidiospores; B. Basidia; C. Caulocystidia; D. Cheilocystidia; E. Pileipellis; F. Stipitipellis. Scale bars = 10 μ m. Drawings by Sana Jabeen.

cell, hyaline in 5% KOH; clamp connections at base common. PLEUROCYSTIDIA absent. CAULOCYSTIDIA at the extreme apex of the stipe, (43.1–)45.2–50.3(–56.11) $\times$ (7.6–)9.4–10.9(–11.2) μm , $\text{avl} \times \text{avw} = 47.7 \times 10.1 \mu\text{m}$, clavate, hyaline in 5% KOH; clamp connections observed at the base. PILEIPELLIS hyphae 4.6–7.7 μm wide, $\text{avw} = 6.1 \mu\text{m}$, septate, filamentous, branched, hyaline in 5% KOH; clamp connections frequent. STIPITPELLIS hyphae (6.1–)8.3–11.04(–11.69) μm wide, $\text{avw} = 9.7 \mu\text{m}$, septate, filamentous, rarely branched; terminal cells clavate; clamp connections observed.

ADDITIONAL SPECIMENS EXAMINED: PAKISTAN. KHYBER PAKHTUNKHWA, Malakand division, Lower Dir district, near Darosh, 1840 m a.s.l., on soil under *Pinus wallichiana*, 4 September 2015, Hira Bashir & Abdul Nasir Khalid D15 (LAH35288; GenBank MG495393); Swat district, Kalam, 2400 m a.s.l., on soil under *C. deodara*, 4 September 2013, Sana Jabeen SJ146 (LAH35045; GenBank MG495395); Swat district, Mashkun, 2500 m a.s.l., on soil under *C. deodara*, 5 September 2013, Aamna Ishaq SJ147 (LAH35046; GenBank MG495396); SJ110 (LAH35289; GenBank MG495394).

HABITAT & DISTRIBUTION—Himalayan moist *Pinaceae*-dominated forests (including *Pinus roxburghii* Sarg., *P. wallichiana*, and few *Cedrus deodara*) and dry oak-dominated temperate forests (including *Quercus oblongata*). Currently known only from Pakistan's Khyber Pakhtunkhwa province.

ECOLOGY & DISTRIBUTION—1840–2500 m asl in dry and moist temperate coniferous forests, Pakistan.

Molecular phylogenetic analysis

A BLAST search of NCBI comparing the ITS consensus sequence of 658 base pairs from the type specimen of *Pseudosperma albobrunneum* (LAH35045) was closely similar to HQ604626 & KP783443 sequences named as *I. sororia* (*P. sororium* (Kauffman) Matheny & Esteve-Rav. in Index Fungorum) from USA and Russia. The ITS dataset involved 100 nucleotide sequences including these two and others closely related from the BLAST along with sequences generated during this study and those from published literature as well as those chosen as outgroup. Of the total 1642 positions in the final dataset, 771 were conserved, 465 were variable, 342 were parsimony informative, and 114 were present as singletons.

In the phylogenetic tree (FIG. 3), two major clades were recovered within *Pseudosperma* labeled as clade A and clade B. The *P. albobrunneum* sequences generated during this study formed their own lineage in clade A clustered with *P. sororium* (HQ604626 & KP783443) but separated from them with 94% boot strap support. The second set of




FIGURE 3. Molecular phylogenetic analysis of *Pseudosperma albobrunneum* based on ITS sequences. Sequences generated during this study are marked with bullets; a red bullet indicates the holotype sequence.

P. sororium sequences (HQ604607, HQ604610, HQ604617, HQ604618) found in clade B does not show remarkable genetic similarity with our *P. albobrunneum* sequences.

Discussion

Pseudosperma albobrunneum is characterized by its whitish to brownish basidiomata, a very prominent long-lasting umbo, a comparatively long and slender stipe, and smooth and elongate to cylindric, sub-reniform basidiospores.

The new Pakistani species resembles the European *P. dulcamaroides* (Kühner) Matheny & Esteve-Rav., which also has brown basidiomata but with a short stipe compared with its cap diameter and with a non-umbonate pileus that becomes convex to flat at maturity (Larsson & al. 2009).

Pseudosperma sororium (Kauffman) Matheny & Esteve-Rav. showed morphological characters more or less similar to *P. albobrunneum* as described by Kauffman (1924), but its yellow pileus and basidiospores that are ellipsoid or elongate-ellipsoid and not truly sub-reniform, sub-inequilateral, obtuse at both ends distinguish *P. sororium* from *P. albobrunneum*. Furthermore, *P. sororium* has a very strong green corn odor.

In the absence of type studies, modern descriptions of *P. dulcamaroides* (Larsson & al. 2009) and *P. sororium* (as *Inocybe*, Stuntz 1978) also support *P. albobrunneum* as morphologically distinct.

Pseudosperma breviterincarnatum (D.E. Stuntz ex Kropp & al.) Matheny & Esteve-Rav. from conifer and quaking aspen forests of the western USA differs by its pinkish lamellae and brownish pileus as well as a pinkish to brownish stipe (Kropp & al. 2013).

Pseudosperma flavorimosum Jabeen & Khalid from Pakistan morphologically resembles *P. albobrunneum* but can be distinguished by its basidiospores that are elliptical, amygdaliform with broad apex and narrow base (Jabeen & Khalid 2020). In the phylogenetic tree (FIG. 3), *P. albobrunneum* forms a separate clade, supporting its novelty.

Acknowledgements

This work was financially supported by Higher Education Commission (HEC)-Pakistan under Indigenous PhD Fellowship (Phase II). Sincere thanks to Dr. Chang-Lin Zhao (Southwest Forestry University, Kunming, Yunnan, P.R. China), Dr. Tine Grebenc (Department of Forest Physiology and Genetics, Slovenian Forestry Institute, Ljubljana, Slovenia), and Dr. Else C. Vellinga (University of California, Berkeley, USA) for presubmission reviews of the manuscript.

Literature cited

- Ahmad S, Iqbal SH, Khalid AN. 1997. Fungi of Pakistan. Sultan Ahmad Mycological Society of Pakistan, Department of Botany, University of the Punjab, Quaid-e-Azam campus, Lahore, 248 p.
- Alvarado P, Manjón JL, Matheny PB, Esteve-Raventós F. 2010. *Tubariomyces*, a new genus of *Inocybaceae*. *Mycologia* 102: 1389–1397. <https://doi.org/10.3852/10-041>
- Bougher NL, Matheny PB. 2011. Two species of *Inocybe* (*Fungi*) introduced into Western Australia. *Nuytsia* 21: 139–148.
- Bougher NL, Matheny PB, Gates GM. 2012. Five new species and records of *Inocybe* (*Agaricales*) from temperate and tropical Australia. *Nuytsia* 22: 57–74.
- Braaten CC, Matheny PB, Viess DL, Wood MG, Williams JH, Bougher NL. 2014. Two new species of *Inocybe* from Australia and North America that include novel secotioid forms. *Botany* 92: 9–22. <https://doi.org/10.1139/cjb-2013-0195>
- Bruns TD. 1995. Thoughts on the processes that maintain local species diversity of ectomycorrhizal fungi. *Plant and Soil* 170(1): 63–73. <https://doi.org/10.1007/BF02183055>
- Champion G, Harry S, Seth K. 1965. Forest types of Pakistan. Pakistan Forest Institute, Peshawar. 233 p.
- Cripps C, Larsson E, Horak E. 2010. Subgenus *Mallocybe* (*Inocybe*) in the Rocky Mountain alpine zone with molecular reference to European Arctic-alpine material. *North American Fungi* 5: 97–126.
- Esteve-Raventós F, Moreno G, Bizio E, Alvarado P. 2015. *Inocybe flavobrunnescens*, a new species in section *Marginatae*. *Mycological Progress* 14: 1–12. <https://doi.org/10.1007/s11557-015-1036-0>
- Fan YG, Bau T. 2013. Two striking *Inocybe* species from Yunnan Province, China. *Mycotaxon*. 123: 169–181. <https://doi.org/10.5248/123.169>
- Fan YG, Bau T. 2014. *Inocybe miyiensis*, a new two-spored species in section *Marginatae* from China. *Nova Hedwigia* 98: 179–185. <https://doi.org/10.1127/0029-5035/2013/0135>
- Farooqi A, Aqduş F, Niazi AR, Jabeen S, Khalid AN. 2017. *Inocybe ahmadii* sp. nov. and a new record of *I. leptocystis* from Pakistan. *Mycotaxon* 132: 257–269. <https://doi.org/10.5248/132.257>
- Gardes M, Bruns TD. 1993. ITS primers with enhanced specificity of basidiomycetes: application to the identification of mycorrhizae and rusts. *Molecular Ecology* 2: 113–118. <https://doi.org/10.1111/j.1365-294X.1993.tb00005.x>
- Hall TA. 1999. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acid Series* 41: 95–98.
- Hamayun M, Khan MA, Begum S. 2003. Marketing of medicinal plants of Utror-Gabral valleys, Swat, Pakistan. *Ethnobotanical Leaflet* 2003: 13.
- Heim R. 1931. Le genre *Inocybe*. *Encyclopédie mycologique* 1. Paris: Paul Lechevalier & Fils. 429 p.
- Horak E. 2005. Röhrlinge und Blätterpilze in Europa. Heidelberg, Germany: Spektrum Akademischer Verlag, 575 p.
- Ilyas S, Razaq A, Khalid AN. 2013. *Inocybe nitidiuscula* and its ectomycorrhizae with *Alnus nitida* from Galyat, Pakistan. *Mycotaxon* 124: 247–254. <https://doi.org/10.5248/124.247>
- Jabeen S, Khalid AN. 2020. *Pseudosperma flavorimosum* sp. nov. from Pakistan. *Mycotaxon* 135: 183–193. <https://doi.org/10.5248/135.183>
- Jabeen S, Ahmad I, Rashid A, Khalid AN. 2016. *Inocybe kohistanensis*, a new species from Swat, Pakistan. *Turkish Journal of Botany* 40: 312–318. <https://doi.org/10.3906/bot-1501-17>

- Kauffman CH. 1924. *Inocybe*. North American Flora 10: 227–261.
- Kobayashi T. 2009. Notes on the genus *Inocybe* of Japan: IV. Species having metuloids collected from Hokkaido, Honshu, and Kyushu. *Mycoscience* 50: 203–211. <https://doi.org/10.1007/S10267-008-0472-Y>
- Kobayashi T, Onishi S. 2010. *Inocybe sericella*, a new species of *Inocybe* sect. *Inocybe* [= *Cortinatae*] from Kobe, Japan. *Nova Hedwigia* 90: 227–232. <https://doi.org/10.1127/0029-5035/2010/0090-0227>
- Kokkonen K, Vauras J. 2012. Eleven new boreal species of *Inocybe* with nodulose spores. *Mycological Progress* 11: 299–341. <https://doi.org/10.1007/s11557011-0783-9>
- Kropp BR, Matheny PB, Hutchison LJ. 2013. *Inocybe* section *Rimosae* in Utah: phylogenetic affinities and new species. *Mycologia* 105: 728–747. <https://doi.org/10.3852/12-185>
- Kühner R, Romagnesi H. 1953. *Flore analytique des champignons supérieurs*. Paris: Masson et Cie. 556 p.
- Kuyper TW. 1986. A revision of the genus *Inocybe* in Europe. 1. Subgenus *Inosperma* and the smooth spored species of subgenus *Inocybe*. *Persoonia Supplement* 3. 247 p.
- Larsson E, Ryberg M, Moreau PA, Mathiesen ÅD, Jacobsson S. 2009. Taxonomy and evolutionary relationships within species of section *Rimosae* (*Inocybe*) based on ITS, LSU and mtSSU sequence data. *Persoonia* 23: 86–98. <https://doi.org/10.3767/003158509X475913>
- Latha KPD, Manimohan P. 2017. *Inocybes* of Kerala. Spore Print Books, Calicut. 181 p.
- Liu L-N, Razaq A, Atri NS, Bau T, Belbahri L, Bouket AC & al. 2018. Fungal systematics and evolution: FUSE 4. *Sydowia* 70: 211–286.
- Massee G. 1904. A monograph of the genus *Inocybe*, Karsten. *Annals of Botany* 18: 459–504. <https://doi.org/10.1093/oxfordjournals.aob.a088972>
- Matheny PB, Bougher NL. 2006. The new genus *Auritella* from Africa and Australia (*Inocybaceae*, *Agaricales*): molecular systematics, taxonomy and historical biogeography. *Mycological Progress* 5: 2–17. <https://doi.org/10.1007/s11557-005-0001-8>
- Matheny PB, Bougher NL. 2017. *Fungi of Australia: Inocybaceae*. Australian Biological Resources Study, Canberra. CSIRO Publishing, Melbourne.
- Matheny PB, Kudzma LV. 2019. New species of *Inocybe* (*Agaricales*) from eastern North America. *Journal of the Torrey Botanical Society* 146: 213–235. <https://doi.org/10.3159/TORREY-D-18-00060.1>
- Matheny PB, Aime MC, Henkel TW. 2003. New species of *Inocybe* from *Dicymbe* forests of Guyana. *Mycological Research* 107(4): 495–505. <https://doi.org/10.1017/S0953756203007627>
- Matheny PB, Aime MC, Bougher NL, Buyck B, Desjardin DE, Horak E, Kropp BR, Lodge DJ, Soyong K, Trappe JM, Hibbett DS. 2009. Out of the Palaeotropics? Historical biogeography and diversification of the cosmopolitan ectomycorrhizal mushroom family *Inocybaceae*. *Journal of Biogeography* 36: 577–592. <https://doi.org/10.1111/j.1365-2699.2008.02055.x>
- Matheny PB, Aime MC, Smith ME, Henkel TW. 2012. Nuevas especies y registros de *Inocybe* (*Agaricales*) de Guyana. *Kurtziana* 37(1): 23–29.
- Matheny PB, Hobbs AM, Esteve-Raventós, F. 2019. Genera of *Inocybaceae*: new skin for the old ceremony. *Mycologia* 112: 83–120. <https://doi.org/10.1080/00275514.2019.1668906>
- Munsell AH. 1975. *Munsell soil color charts*. Macbeth Division of Kollmorgen Corporation. Baltimore, Maryland.
- Naseer A, Khalid AN, Smith ME. 2017. *Inocybe shawarensis* sp. nov. in the *Inosperma* clade from Pakistan. *Mycotaxon* 132(4): 909–918. <https://doi.org/10.5248/132.909>
- Naseer A, Ghani S, Niazi AR, Khalid AN. 2019. *Inocybe caroticolor* from oak forests of Pakistan. *Mycotaxon* 134(2): 241–251. <https://doi.org/10.5248/134.241>

- Ryberg M, Larsson E, Jacobsson S. 2010. An evolutionary perspective on morphological and ecological characters in the mushroom family *Inocybaceae* (*Agaricomycotina*, *Fungi*). *Molecular Phylogenetics and Evolution* 55: 431–442. <https://doi.org/10.1016/j.ympev.2010.02.011>
- Saba M, Ahmad I, Khalid AN. 2015. New reports of *Inocybe* from pine forests in Pakistan. *Mycotaxon* 130: 671–681. <https://doi.org/10.5248/130.671>
- Saba M, Haelewaters D, Pfister DH, Khalid AN. 2020. New species of *Pseudosperma* (*Agaricales*, *Inocybaceae*) from Pakistan revealed by morphology and multi-locus phylogenetic reconstruction. *MycKeys* 69: 1–31. <https://doi.org/10.3897/mycokeys.69.33563>
- Siddiqui MF, Shaukat SS, Ahmed M, Khan N, Khan IA. 2013. Vegetation-environment relationship of conifer dominating forests of moist temperate belt of Himalayan and Hindukush regions of Pakistan. *Pakistan Journal of Botany* 45: 577–92.
- Singer R. 1986. *The Agaricales in modern taxonomy*. 4th ed. Koenigstein, Germany: Koeltz Scientific Books. 981 p.
- Stucki B, Khan HA. 1999. Working plan for Utror-Desan forests of Kalam Forest Division. Peshawar, Pakistan Nizam Printing Press.
- Stuntz D. 1978. Interim 'skeleton key' to some common species of *Inocybe* in the Pacific Northwest. *Pacific Northwest Key Council* (rev. ed. 2004). Accessed 30 December 2020: <https://www.svims.ca/council/Inocyb.htm>
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30: 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Ullah Z, Jabeen S, Ahmad H, Khalid AN. 2018. *Inocybe pakistanensis*, a new species in section *Rimosae*. *Phytotaxa* 348(4): 279–288. <https://doi.org/10.11646/phytotaxa.348.4.4>
- White TJ, Bruns TD, Lee SB, Taylor JW. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. 315–322, in: MA Innis & al. (eds). *PCR protocols: a guide to methods applications*. Academic Press, New York. <https://doi.org/10.1016/B978-0-12-372180-8.50042-1>